

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 0 972 838 A1

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:  
19.01.2000 Bulletin 2000/03

(21) Application number: 99113182.2

(22) Date of filing: 08.07.1999

(51) Int. Cl.<sup>7</sup>: C12N 15/70, C12N 15/64,  
C12N 15/65, C07K 14/56  
// C12N15/21

(84) Designated Contracting States:  
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE  
Designated Extension States:  
AL LT LV MK RO SI

(30) Priority: 15.07.1998 EP 98113156  
09.10.1998 EP 98119078

(71) Applicant:  
Roche Diagnostics GmbH  
68305 Mannheim (DE)


(72) Inventors:  
• Kopetzki, Erhard  
82377 Penzberg (DE)  
• Schantz, Christian  
85402 Thalhausen (DE)

(74) Representative:  
Notegen, Eric-André, Dr.  
Grenzacherstrasse 124  
4070 Basel (CH)

(54) **Escherichia coli host/vector system based on antibiotic-free selection by complementation of an auxotrophy**

(57) A minimal prokaryotic expression vector which cannot be homologously recombined with the genome of prokaryotic organisms containing: e) an origin of replication, f) an auxotrophy marker gene g) a promoter which is functionally active in prokaryotes and h) a for-

eign gene to be expressed which is under the control of the said promoter is particularly suitable for antibiotic-free selection.



EP 0 972 838 A1

## Description

[0001] The invention concerns new prokaryotic expression systems which enable an antibiotic-free selection and their use for the production of recombinant proteins.

[0002] Known prokaryotic expression plasmids contain, in addition to one or several antibiotic resistance gene(s), additional DNA sequences which are not required and burden the cell metabolism. These include DNA sequences resulting from the cloning process, DNA segments that are relics from multipurpose vectors such as e.g. specific promoters for the in vitro synthesis of mRNA and specific phage replication origins for the synthesis of single stranded DNA and extend to rudimentary duplicated vector sequences which can be the cause of undesired plasmid rearrangements.

[0003] The presence of a plasmid and in particular of an expression vector causes an additional metabolic burden for the cell. This results in a selection pressure which favours the formation of cells without plasmids which can be largely prevented by antibiotic selection.

[0004] Antibiotics such as ampicillin, tetracycline, kanamycin and chloramphenicol are usually used for the selection. Especially the use of  $\beta$ -lactam antibiotics such as ampicillin is problematic in the production (fermentation) of therapeutic products.

[0005] For the reasons discussed above antibiotic plasmid selection by means of  $\beta$ -lactam antibiotics is not used in recent methods. In some cases the host/vector system that was used proved to be so stable that it was possible to omit a plasmid selection during the preculture and during the main fermentation (Weir, A.N.C. and Mountain, A., EP 0651803). However, as a rule this is associated with a reduced product yield. In cases in which antibiotic selection is indispensable, tetracycline is often used as an alternative to ampicillin (Carter, P. et al., 1992). In contrast to the  $\beta$ -lactam antibiotics, the tetracyclines are not reactive chemical compounds and cannot be inactivated by enzymatic modification during fermentation. The tetracycline resistance gene codes for a protein which modifies the bacterial membrane and thus prevents the antibiotic from entering the cell.

[0006] Struhl, K. and coworkers (Struhl, K. et al., 1976) using imidazole glycerol phosphate dehydratase (HIS3) as an example, show that an appropriate *E. coli* mutant (hisB) can be directly complemented by means of plasmids containing yeast DNA and that the yeast enzyme coded by HIS3 is functionally expressed in *E. coli*. This ability of genes to complement mutations in *E. coli* was used as a selection criterion to clone for example (complementation cloning) other yeast genes (LEU2, URA3, TRP5 and ARG4).

[0007] *E. coli* host strains with a stable mutation (reversion  $>10^{-10}$ ; preferably non-revertable deletion mutants) are required for the selection by complementation. However, the reversion rate of known mutations is of the order of magnitude of  $<10^{-10}$  (Schlegel, H.G., 1981).

[0008] The known *E. coli* laboratory strains differ with regard to their genotype in individual mutations which in many cases were produced by undirected mutagenesis by radiation (X-ray or UV radiation), chemical mutagens (e.g. base analogues, nitrous acid and alkylating compounds) or biological techniques (e.g. phage Mu and transposon mutagenesis (Bachman, B.J., 1986).

[0009] The object of the invention was to develop a stable prokaryotic (preferably *E. coli*) host/vector system (expression system) which avoids a plasmid selection by antibiotics. The selection principle is based on complementation of a stable auxotrophy of the prokaryotic host cell by an adequate yeast gene.

[0010] The invention concerns a minimal prokaryotic expression vector which cannot be homologously recombined with the genome of prokaryotic organisms containing:

- a) an origin of replication,
- b) an auxotrophy marker gene,
- c) a promoter which is functionally active in prokaryotes and
- d) a foreign gene (foreign sequence) to be expressed under the control of the said promoter.

[0011] In this connection the transcription of the auxotrophy marker gene and the transcription of the foreign gene to be expressed are preferably in opposite directions. It is also preferred that the marker gene is heterologous to the prokaryotic organism used as the host cell. It is additionally preferred that the expression of the auxotrophy marker gene in the host/vector system described in the invention is 1 % or less of the total protein expressed in the host cell when the auxotrophy is complemented in the host cell.

[0012] In a preferred embodiment the expression vector according to the invention is characterized in that the reading frame of the auxotrophy marker gene and of the gene to be expressed are oppositely oriented and are terminated by the same transcription terminator or several transcription terminators arranged in series.

[0013] An auxotrophy marker gene is understood as a gene which enables cell growth of auxotrophic host strains under selective culture conditions. Particularly preferred marker genes are heterologous eukaryotic genes, preferably yeast genes, which compensate a defect (mutation) in an essential metabolic path of the host organism. With *E. coli* as the host organism, mutants with a defect in the tryptophan, leucine or uracil biosynthetic pathway are preferred which

can then be complemented by the marker gene.

[0014] A further subject matter of the invention is a prokaryotic auxotrophic host cell which contains a vector according to the invention wherein this host cell has a mutation (deletion) in a gene which is complemented by the selection marker gene of the expression vector and the deletion or mutation has the effect that no functional product of the said host cell gene is expressed. The deletion is preferably located in an essential chromosomal gene which corresponds to the auxotrophy marker gene of the expression vector. A further subject matter of the invention is a method for producing a prokaryotic expression system which comprises introducing an expression vector according to the invention into a prokaryotic auxotrophic cell with a deletion in a gene which corresponds to the auxotrophy marker of the expression vector where the deletion or mutation has the effect that no functional product of the said host cell gene is expressed.

[0015] A further subject matter of the invention is a process for the production of a heterologous protein by heterologous expression of a desired protein by using an expression vector according to the invention in a host cell according to the invention and isolation of the expression product from the cells or the supernatant wherein the fermentation is carried out in a minimal medium or in a synthetic complete medium.

[0016] A further subject matter of the invention is a process for the production of a prokaryotic expression system which is characterized in that an expression vector according to the invention is introduced into a prokaryotic host cell with a mutation in a chromosomal gene which is complemented by the auxotrophy marker gene of the expression vector and the chromosomal mutation has the effect that no functional product of the said chromosomal host cell gene can be expressed.

[0017] In the present invention a new *E. coli* host/vector system based on the complementation of an essential chromosomal *E. coli* auxotrophy (*trpC*, *pyrF*) by the plasmid-coded expression of a suitable yeast gene (*TRP1*, *URA3*) was developed instead of the usual selection with antibiotics. In addition:

- i) new expression vectors optimized (minimized) with respect to size were constructed using minimal functional elements (selection marker, origin of replication and expression cassette),
- ii) non-revertable host strains were isolated by homologous recombination (deletion) of a desired chromosomal *E. coli* gene (*trpC*, *pyrF*) and
- iii) the components of the new host/vector system (expression plasmids and host strains) were checked with regard to function and performance on the basis of the expression of a model gene (interferon- $\alpha$ -2b).

[0018] For this:

- i) *E. coli* host strains with stable mutations (deletions) of the chromosomal genes *trpC* and *pyrF* and
- ii) vectors which contain the complementing yeast genes *TRP1* (Tschumper, G., 1980) or *URA3* (Rose, M., 1984) were constructed.

[0019] Double mutants or deletion mutants are preferably constructed as stable host strains.

[0020] The minimal vectors according to the invention are preferably very small (smaller than 2000 bp) and only contain the absolutely essential elements of an expression plasmid i.e. an origin of replication, a heterologous selection marker gene and a heterologous expression cassette and these have the smallest possible size to avoid homologous recombination with the genome of the prokaryotic host cell. A homologous recombination within the meaning of the invention is understood as the exchange of DNA sequences between the vector according to the invention and the genome of the host cell. Homologous recombination can occur when there is a considerable sequence compatibility (at least 50 - 1000 bp) between the vector and genome. Less agreement in the sequence (below 50 bp) does not lead to homologous recombination to a detectable extent within the meaning of the invention.

[0021] Examples of suitable host cells are *E. coli* *TrpC* mutants which are characterized by an inactive N-(5'-phosphoribosyl)-anthranilate isomerase (EC 5.3.1.24) and *pyrF* mutants which are characterized by an inactive orotidine-5'-phosphate decarboxylase (EC 4.1.1.23). The mutants are incapable of synthesizing tryptophan or uracil and can therefore not grow on nutrient media which lack these components. In contrast they exhibit normal growth on complete medium since tryptophan and uracil are present in adequate amounts.

[0022] The auxotrophic *trpC* and *pyrF* mutants are complemented by a plasmid according to the invention which contains the absent or defective gene for the respective biosynthesis without a mutation. However, the complementing genes are not derived from *E. coli* but from the yeast *Saccharomyces cerevisiae*. They code for the same enzymes as the corresponding *E. coli* genes. Due to the poor function of the yeast promoters, their transcription in *E. coli* is less than the transcription of the host's endogenous analogous *E. coli* genes. The plasmid-coded expression of the *URA3* yeast gene (*pBR*<sub>322</sub> plasmid derivative) is for example only one third of the expression of the analogous chromosomal *pyrF* gene (Rose, M., 1984). This has the advantage that the marker genes are not overexpressed despite the gene dose effect and the cell is not additionally metabolically stressed.

[0023] Plasmids are extrachromosomal (episomal) circular double-stranded DNA molecules which have a length of

several thousand nucleotide building blocks and occur in several to several hundred copies per cell. They have a DNA segment, the so-called origin of replication, which enables an autonomous plasmid replication/amplification independent of the chromosomal DNA replication.

[0024] A stable E. coli host/vector system was developed which avoids plasmid selection by antibiotics. The selection principle is based on the complementation of a stable E. coli auxotrophy by an adequate yeast gene.

[0025] The E. coli host strains preferably contain a stable, preferably non-revertible mutation (deletion) in the chromosomal genes *trpC* and/or *pyrF* and an expression plasmid which contains the complementing yeast genes *TRP1* and/or *URA3*.

[0026] The basic expression plasmids are characterized by:

- i) the absence of antibiotic resistance genes and
- ii) their very small size (< 2000 bp).

[0027] They only contain the absolutely essential elements for replication in E. coli and expression of a desired gene:

- i) an origin of replication,
- ii) a selection marker gene and
- iii) a prokaryotic or eukaryotic expression cassette (gene therapy), and the size of these plasmid elements is as minimal as possible.

## The origin of replication

[0028] The origin of replication for the autonomous replication of the plasmid in the cell is derived from a prokaryotic plasmid, preferably from a natural E. coli plasmid. Most of the E. coli plasmids used today are derivatives of the cloning vector pBR<sub>322</sub> (Boliver, F. et al., 1979) the replicon of which (origin of DNA replication) is based on the plasmid pMB1 (Sutcliffe, G. et al. 1979) which only differs slightly from the plasmid ColE1. The replication of the plasmid pBR<sub>322</sub> is also referred to as replication of the ColE1 type (Backman, K. et al., 1978). The minimal DNA sequence which is responsible for an autonomous replication of the pBR<sub>322</sub> plasmid was reduced by Backman, K. et al. 1978 to a 580-650 bp long DNA sequence. The plasmid copy number per cell is determined by the type of replication origin. Depending on the number of copies per cell that can be achieved, one refers to low copy number (ca. 10 copies, pACYC plasmids), medium copy number (ca. 50 copies, pBR plasmids) and high copy number (ca. 500 copies, pUC plasmids) plasmids. The high copy number origin of replication of the pUC plasmids is only based on a point mutation of the pBR origin of replication (Chambers, S.P. et al., 1988). The expression yield of a gene generally correlates with the copy number of the gene to be expressed.

## The expression cassette

[0029] An expression cassette (heterologous transcription unit) is composed of:

- i) a promoter,
- ii) a ribosomal binding site,
- iii) a multiple cloning cleavage site and
- iv) a transcription terminator (rho-independent termination).

[0030] The multiple cloning cleavage site serves to insert the gene (structural gene) to be expressed and for this purpose it is composed of at least two restriction endonuclease cleavage sites that are present only once in the vector. Cleavage sites are preferably used for the cleavage site that faces the promoter which contain the nucleotide sequence for an ATG start codon in their recognition sequence such as e.g. the recognition sequences for the restriction enzymes *NcoI*, *BspHI*, *SpHI* and *NdeI*. This facilitates the precise connection between promoter and structural gene.

[0031] A promoter is a DNA fragment which contains the control signals for the start of mRNA synthesis and the frequency with which this mRNA molecule is to be formed and is functionally active in an expression cassette. A typical natural E. coli promoter is 80 - 300 base pairs long and contains several characteristic (homologous) regions which are common to the various promoters and which have been determined by sequence comparisons (Rosenberg, M. and Court, D., 1979; Harley, C.B. and Reynolds, R.P., 1987). The highly conserved regions are:

- i) the recognition site for the RNA polymerase with the motif 5'-TTGACA-3', which is 35 base pairs before the start of the mRNA synthesis,
- ii) the Pribnow Schaller box or so-called TATA box with the codonical prototype sequence 5'-TATAAT-3' which is

located 10 base pairs before the start of the mRNA synthesis and  
 iii) an AT-rich region around position "-43" in strong promoters.

[0032] Depending on the frequency with which a promoter is read one refers to weak and strong promoters. A distinction is also made between constitutive and regulatable (inducible, derepressible) promoters. A good promoter in terms of the industrial production of proteins is usually a strong, strictly regulated promoter that is inactive during cell multiplication and which can be specifically switched on as required e.g. at the end of a fermentation. These requirements of a very high and at the same time strictly controllable promoter activity have led to the construction of the hybrid promoters that are nowadays preferably used. In the case of the tac hybrid promoter which can be readily regulated, the "-35" region of the strong constitutive trp promoter was fused with a "-10" region of the inducible lac promoter (DeBoer, H.A. et al., 1983; Amann, E. et al., 1983). A further example is the T5-P<sub>N25/03/04</sub> hybrid promoter (Stüber, D. et al., 1990) which is based on a strong constitutive T5 bacteriophage promoter and was reconstructed by combination/insertion of two lac operators (lacO) to form a strong promoter that can be readily regulated. The tac as well as the T5-P<sub>N25/03/04</sub> hybrid promoter are negatively regulated by the lacI repressor which prevents transcription of the promoter by interaction with the lac operator(s). Addition of the natural inducer lactose or of the very potent non-metabolizable lactose analogue isopropyl-β-D-thiogalactopyranoside (IPTG) leads to detachment of the lacI repressor and thus to formation of the desired mRNA and protein synthesis.

[0033] Transcription terminators are DNA sequences of 50-100 base pairs in length which give the RNA polymerase the signal for termination of the mRNA synthesis. They are structurally characterized by formation of a hairpin-shaped secondary structure resulting from complementary base pairing (mostly G/C-rich). Functionally they act independently of the orientation. Very efficient (strong) terminators at the 3' end of an expression cassette are advisable to prevent the RNA polymerase from reading through particularly when using strong promoters. Inefficient transcription terminators can lead to the formation of an operon-like mRNA which can be the reason for an undesired plasmid-coded gene expression. Furthermore the secondary structure at the 3' end of the mRNA caused by the terminator impedes degradation by 3' exonucleases (Higgins, C.F. et al., 1993) which can increase the half-life of the mRNA and consequently also increase the amount of synthesized protein. Examples of often used transcription terminators are the terminator T0 of the bacteriophage λ (Schwarz, E. et al., 1978), the fd bacteriophage terminator (Beck E. and Zink, B., 1981) and the terminator T1 of the E. coli rrnB operon. (Brosius, J. et al., 1981).

[0034] A ribosomal binding site (RBS) is also required for efficient initiation of protein biosynthesis. It comprises the sequence from the ATG start codon up to the co-called Shine Dalgarno sequence with the consensus motif AGGA. The distance between the ATG start codon and the SD sequence varies between 5-15 base pairs and is A/T-rich. The distance between the starting point of mRNA synthesis and the ATG translation start codon varies as a rule between 15 - 80 base pairs.

[0035] Any sequences are suitable as a foreign sequence within the sense of the invention. Such nucleic acid sequences preferably code for polypeptides which have an effect on metabolism in the human body. Such proteins or polypeptides are preferably therapeutically relevant polypeptides such as cytokines, proteins or protein hormones.

[0036] There are a large number of techniques available for producing E. coli mutants which are suitable for the construction of E. coli host strains. They range from mutagenesis of the whole cell followed by selection of the desired phenotype to the specific mutagenesis of individual bases or of exactly defined gene or genome sections. The methods are described by Miller, J.H. (1972), Neidhardt, F.C. et al. (1987) and Winnacker, E.-L. (1985).

[0037] A stable non-revertible auxotrophy marker was introduced into host strains by the directed deletion of the complete gene or of a large gene fragment by means of homologous recombination according to the method of Hamilton C.M. et al., 1989. The original genotype is preserved in this technique i.e. no additional mutations are introduced into the E. coli genome. The principle of the gene exchange technique is elucidated in more detail using the construction of a tryC deletion mutant as an example.

[0038] It is intended for example to delete the complete trpC gene (1355 bp) from the tryptophan operon of E. coli (Yanofsky, C. et al., 1981). For this the flanking regions (700 - 800 bp in each case) of the trpC gene are amplified by means of PCR using appropriate primers and chromosomal E. coli DNA as the template. The overhangs of the primers contain singular cleavage sites for restriction enzymes in order to ligate the amplified DNA fragments of the genes trpD and trpB into the vector pMAK705 in a direct fusion. The vector pMAK705 is characterized by a temperature-sensitive origin of replication (origin of replication from pSC101) which is inactive at 44°C and by a chloramphenicol resistance gene. In order to delete the chromosomal trpC gene, the pMAK705-trpD/B plasmid is transformed into the desired E. coli strain. A prerequisite for a homologous recombination between the plasmid and E. coli chromosome is a recombination-competent E. coli strain (recA<sup>+</sup> genotype). Strains with a recA<sup>-</sup> genotype cannot carry out a homologous recombination and are therefore unsuitable. The transformed cells are cultured at 44°C in the presence of chloramphenicol. Since the plasmid does not replicate at 44°C due to its temperature-sensitive origin of replication (ori<sup>ts</sup>), only those cells survive which have integrated the vector into the chromosome for example by means of homologous recombination. The co-integrates are isolated and cultured under chloramphenicol selection over several passages at 30°C. The chro-

mosomally localized origin of replication of the plasmid pMAK705-trpD/B which is active at this temperature leads to a drastic reduction in the growth rate of the co-integrates. Cells which have removed the plasmid from the chromosome by a second recombination event grow normally again. They contain the plasmid detached from the chromosome which replicates at 30°C in the presence of chloramphenicol. The result is that cells containing plasmids overgrow the co-integrates and dominate in the culture after several passages. As elucidated in Figure 1, the integrated plasmid can be detached from the chromosome in two different ways A and B. The plasmid then contains the gene trpC (B) to be deleted or it is again present in its original form in the cell (A) depending on the homologous recombination event. Cells which contain a plasmid with the chromosomal trpC gene very probably have the desired trpC deletion. They are identified by restriction analysis of the plasmids and are cured of the plasmid by a single culture at 44°C and several cultures at 30°C in the absence of chloramphenicol. Subsequently the clones are tested with regard to trpC auxotrophy by plating out the individual colonies on a medium with and without tryptophan (replica plating).

#### Special E. coli host/vector systems and host strains

[0039] An extremely strictly regulated host/vector system was developed especially for the expression of proteins that are toxic for E. coli in order to prevent a basal expression of a gene which may still cause interference. The T7 polymerase system of Studier, F. et al., 1990 is based on a special RNA polymerase from the bacteriophage T7 which is absolutely selective and specific for the T7 promoter. Since the host's E. coli RNA polymerase does not recognize this T7 promoter, the T7 polymerase is provided at the time of induction by

- i) infecting the culture with a suitable phage such as e.g. the  $\lambda$  bacteriophage derivative CE6 or
- ii) by inducing a chromosomally-integrated T7 polymerase under lacUV5/lacI promoter control.

[0040] The special E. coli host strains BL21(DE3) and HMS174(DE3) containing a T7 polymerase gene were constructed by chromosomal insertion of the lysogenic  $\lambda$  bacteriophage derivative DE3. This system which is often used in research laboratories is less suitable for industrial purposes because:

- i) only a limited number of host strains (2) are available,
- ii) the DE3 lysogen is not absolutely stable in a fermenter and
- iii) the growth of the cells stagnates after the induction and consequently a growth to high cell densities is difficult.

[0041] The following examples, publications, sequence protocols and figures further elucidate the invention the protective scope of which results from the patent claims. The described methods are to be understood as examples which still describe the subject matter of the invention even after modifications.

The figures show:

[0042]

**Figure 1:** Construction of E. coli trpC deletion mutants by a gene exchange technique. The sequences flanking the trpC gene to be deleted are integrated into the vector pMAK705. Only E. coli strains transformed by homologous recombination and integration of the plasmid into the chromosome and cultured at 44°C with the addition of chloramphenicol survive since the plasmid cannot replicate (co-integrate formation) at 44°C. Subsequent culture at 30°C under chloramphenicol selection leads to a second homologous recombination in which the plasmid is cut out of the chromosome. This break up of the co-integrate can lead to the initial plasmid pMAK705-trpD/B (A) or to a pMAK705-trpD/C/B plasmid derivative which additionally contains the trpC gene (B). The trpC gene on the E. coli chromosome is only deleted in case B.

**Figure 2:** Construction of the base vectors OripBR-TRP1, OripBR-URA3 and OripBR-TET: the plasmids OripBR-TRP1 (1497 bp), OripBR-URA3 (1697 bp) and OripBR-TET (2012 bp) are formed utilizing the cleavage sites SfiI and XhoI by ligating the marker genes TET, URA3 and TRP1 isolated from the plasmids pBR<sub>322</sub>, yEP24 and yRP7 with a DNA segment containing the origin of replication from pBR<sub>322</sub>.

**Figure 3:** Nucleotide sequence of the expression cassette composed of the T5-P<sub>N25/03/04</sub> hybrid promoter, the ribosomal binding site RBSII, a multiple cloning cleavage site (NdeI/SalI/XmaI/EcoRV/CelII) and the  $\lambda$ -T0 terminator.

**General methods:**

[0043] Standard methods were used to manipulate DNA as described in Sambrook, J. et al. (1989) In: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, Cold, Spring Harbor, New York. The molecular biological reagents were used according to the manufacturer's instructions.

[0044] The origin of replication from the vector pBR<sub>322</sub> (medium copy number) and from the vector pUC20 (high copy number) were used as the **origin of replication**. The minimal DNA sequence for the autonomous replication of these plasmids is between 580 - 650 bp (Backman, K. et al., 1978).

[0045] The yeast genes TRP1 and URA3 were selected as selection marker genes. The tetracycline resistance gene was used to construct isogenic standard reference plasmids.

**Example 1****Construction of the base vectors OripBR-TRP1, OripBR-URA3 and OripBR-TET****1.1 Construction of the base vector OripBR-TRP1**

[0046] The plasmid OripBR-TRP1 was composed of two DNA fragments which were obtained by the polymerase chain reaction (PCR) according to the method of Mullis, K.B. and Faloona, F.A., 1987. In a first PCR reaction the origin of replication of the plasmid pBR<sub>322</sub> from bp position 2517 - 3160 according to the publication of Sutcliffe J.G., 1979 was amplified using the primers N1 (SEQ ID NO:1) and N2 (SEQ ID NO:2).

SfiI

N1: 5' -AAAAAAGGCCATATAGGCCGCGCGTTGCTGGCGTTTTTC-3'

XhoI

N2: 5' -AAAAAACTCGAGGAGCGTCAGACCCCGTAGAAAAG-3'

and pBR<sub>322</sub> as the template DNA. A singular SfiI restriction endonuclease cleavage site was introduced by means of the 5' overhanging end of the PCR primer N1 and a singular XhoI restriction endonuclease cleavage site was introduced by means of the 5' overhanging end of the PCR primer N2.

[0047] The TRP1 yeast gene (5' flanking region and the TRP1 structural gene) from bp position 22 - 777 according to the publication of Tschumper, G. and Carbon, J., 1980 was amplified in a second PCR reaction using the primers N3 (SEQ ID NO:3) and N4 (SEQ ID NO:4)

SfiI

NotI

N3: 5' -AAAAAAGGCCATATAGGCCATATGCGGCCGCGAGAGGGCCAAGAGGGAGGGC-3'

XhoI

BamHI

fd-terminator

N4: 5' -AAAAAACTCGAGATATGGATCCTCTCCAAAAAAAAGGCTCCAAAAGGAGCCT

TTAATTGTATCGGTTTATTACTATTTCTTAGCATTTTTGACGAAATTTGC-3'

EndEnd

and YRP7 (Kopetzki, E. et al., 1989) as the template DNA. A singular SfiI cleavage site and NotI cleavage site were introduced by means of the 5' overhanging end of the PCR primer N3 and a second translation stop codon, the fd transcription terminator of the bacteriophage fd from position 1522-1570 (Beck, E. and Zink, B., 1981) and two singular restriction endonuclease cleavage sites (MAKHI and XhoI) were introduced at the 3' end of the coding region of the TRP1 structural gene by means of the 5' overhanging end of the PCR primer N4.

[0048] The two PCR fragments digested with SfiI and XhoI were ligated after purification by agarose gel electrophoresis. Afterwards the E. coli K12 strain JA300 (thr, leuB6, thi, thyA, trpC1117, hsrK, HsmK, strR; DSMZ 4776) (source: "Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH) was transformed by means of calcium chloride according to the method of Hanahan, D., 1983 or by electroporation according to the protocol of Fiedler, S. and Wirth, R., 1988. The transformants were selected on agar plates in M9 minimal medium (preparation: see Sambrook, J. et al., 1989) and 0.5 % (w/v) casamino acids (acid-hydrolysed casein contains no tryptophan) from the Difco Company and subsequently grown in liquid culture. The desired plasmid OripBR-TRP1 (1497 bp) was identified by restriction mapping.

## 1.2 Construction of the base vector OripBR-URA3

[0049] The plasmid OripBR-URA3 was composed of two PCR fragments like the plasmid OripBR-TRP1. However, the URA3 yeast gene was used instead of the TRP1 selection marker gene. First PCR reaction: see example 1.1.

[0050] The URA3 yeast gene (5' flanking region and the URA3 structural gene) from bp position 75 - 1030 according to the publication of Rose, M., 1984 was amplified in a second PCR reaction using the primers N5 (SEQ ID NO:5) and N6 (SEQ ID NO:6)

SfiI                      NotI

N5: 5' -AAAAAAGGCCATATAGGCCATATGCGGCCGCCGTAATCTCCGAACAGAAGGAAGA  
AC-3'

XhoI              BamHI                      fd-terminator

N6: 5' -AAAAAAGCTCGAGACGAGGATCCATAAACCGATACAATTAAGGCTCCTTTTGGAGC  
CTTTTCTTTTGGAGAACTTTAGTTTGTGCGCCGCATCTTCTCAA-3'  
EndEnd

and YEp24 (Botstein, D. et al., 1979) as the template DNA. A singular SfiI cleavage site and NotI cleavage site were introduced by means of the 5' overhanging end of the PCR primer N5 and a second translation stop codon, the fd transcription terminator of the bacteriophage fd (anti-native orientation) from position 1522-1570 (Beck, E. and Zink, B., 1981) and two singular restriction endonuclease cleavage sites (BamHI and XhoI) were introduced at the 3' end of the coding region of the TRP1 structural gene by means of the 5' overhanging end of the PCR primer N6.

[0051] The two PCR fragments digested with SfiI and XhoI were ligated after purification by agarose gel electrophoresis. Afterwards the E. coli K12 strain CSH28 ( $\Delta$ lacpro, supF, trp, pyrF, his, strA, thi) (source: Cold Spring Harbor Laboratory, New York; Miller, J.H., 1972) was transformed by means of calcium chloride according to the method of Hanahan, D., 1983 or by electroporation according to the protocol of Fiedler, S. and Wirth, R., 1988. The transformants were selected on agar plates in M9 minimal medium (preparation: see Sambrook, J. et al., 1989) containing 0.5 % (w/v) casamino acids from the Difco Company and 5 mg/ml tryptophan and subsequently grown in liquid culture. The desired plasmid OripBR-URA3 (1697 bp) was identified by restriction mapping.

## 1.3 Construction of the antibiotic-resistance-containing reference vector OripBR-TET

[0052] The plasmid OripBR-TET was composed of two PCR fragments like the plasmids OripBR-TRP1 and OripBR-URA3. However, the tetracycline resistance gene (TET) from the plasmid pBR<sub>322</sub> was used instead of the TRP1 or URA3 selection marker gene. First PCR reaction: see example 1.1.

[0053] The TET resistance gene (promoter and the TET structural gene) from bp position 3 - 1275 according to the publication of Sutcliffe J.G., 1979 was amplified in a second PCR reaction using the primers N7 (SEQ ID NO:7) and N8 (SEQ ID NO:8)



SfiI                      NotI

5        N7: 5' -AAAAAAGGCCATATAGGCCATATGCGGCCGCTCATGTTTGACAGCTTATCATCGA  
              TAAG-3'

XhoI              XbaI                      fd terminator

10       N8: 5' -AAAAAACTCGAGATATTCTAGAATAAACCGATACAATTAAAGGCTCCTTTTGGAGC  
              CTTTTTTTTTTGGAGATCAGGTCGAGGTGGCCCGGCTCC-3'

15       and pBR<sub>322</sub> (Sutcliffe, J.G., 1979) as the template DNA. A singular SfiI cleavage site and NotI cleavage site were introduced by means of the 5' overhanging end of the PCR primer N7 and the fd transcription terminator of the bacteriophage fd (anti-native orientation) from position 1522-1570 (Beck, E. and Zink, B., 1981) and two singular restriction endonuclease cleavage sites (XhoI and XbaI) were introduced at the 3' end of the coding region of the TET structural gene by means of the 5' overhanging end of the PCR primer N8.

20       [0054] The two PCR fragments digested with SfiI and XhoI were ligated after purification by agarose gel electrophoresis. Afterwards the E. coli K12 strain JA300 (thr, leuB6, thi, thyA, trpC1117, hsrK, HsmK, strR; DSMZ 4776) (source: "Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH) was transformed by means of calcium chloride according to the method of Hanahan, D., 1983 or by electroporation according the protocol of Fiedler, S. and Wirth, R.,  
 25       1988. The transformants were selected (agar plates) in LB complete medium (preparation: see Sambrook, J. et al., 1989) containing 12.5 µg/ml tetracycline and subsequently grown in liquid culture. The desired plasmid OripBR-TET (2012 bp) was identified by restriction mapping.

#### 1.4 Removal of an NdeI cleavage site in the plasmids OripBR-TRP1, OripBR-URA3 and OripBR-TET

30       [0055] An undesired NdeI cleavage site which was formed from the 2 PCR fragments between the cleavage sites NotI and SfiI during ligation of the base vectors OripBR-TRP1, OripBR-URA3 and OripBR-TET, was subsequently removed by means of an NotI/SfiI nucleotide adapter.

35       [0056] For this the plasmids OripBR-TRP1, OripBR-URA3 and OripBR-TET were digested with NotI and SfiI, the vector fragments were purified by agarose gel electrophoresis and ligated with an NotI/SfiI nucleotide adapter. The desired plasmids were identified by restriction mapping (missing NdeI cleavage site).

40       [0057] The NotI/SfiI adapter was prepared by hybridization from the mutually complementary oligonucleotides N9 (SEQ ID NO:9) and N10 (SEQ ID NO:10). For this 10 nmol of each of the oligonucleotides N9 and N10 were mixed in 100 µl 12.5 mmol/l Tris-HCl, pH 7.0 and 12.5 mmol/l MgCl<sub>2</sub>, the mixture was heated for 5 min to 95°C and subsequently slowly cooled to room temperature.

N9 : 5' -AGGCCGGGGGGGGGC-3'

N10 : 5' -GGCCGCCCCCCCCCGCCTATA-3'

5

3-ATATCCGGCCCCCCCCCGCCGG-5'

Not I

10

## 15 Example 2

### Construction of the prokaryotic expression cassette

20 **[0059]** The expression cassette that was used (heterologous transcription unit) was composed of the following elements:

- 25 i) the regulated T5-P<sub>N25/03/04</sub> hybrid promoter (Bujard, H. et al, 1987; Stüber, D. et al, 1990),  
ii) a synthetic ribosomal binding site RBSII (Stüber, D. et al., 1990),  
iii) a multiple cloning cleavage site with the singular cleavage sites NdeI, SalI, XmaI, EcoRV and CclII and  
iv) the  $\lambda$ -T0 bacteriophage transcription terminator (Schwarz, E. et al., 1978).

**[0060]** The T5-P<sub>N25/03/04</sub> hybrid promoter (Fig. 3: bp position: 1 - 87) was linked to the  $\lambda$ -T0 terminator (Fig. 3: bp position: 146 - 241) by means of an EcoRI-RBSII-NdeI/ Sall/XmaI/EcoRV/CelII linker.

**[0061]** The expression cassette was obtained by reconstruction of a pDS56/RBSII plasmid (Stüber, D. et al., 1990).  
30 For this a derivative of the pDS56/RBSII plasmid was digested with EcoRI and CclII and the EcoRI/CclII-pDS56 vector fragment purified by agarose gel electrophoresis was ligated with an EcoRI-RBSII-NdeI/SalI/XmaI/EcoRV/ CclII linker.

**[0062]** The EcoRI-RBSII-NdeI/Sall/XmaI/EcoRV/ CeliI linker was prepared by hybridization from the mutually complementary oligonucleotides N11 (SEQ ID NO:11) and N12 (SEQ ID NO:12). 10 nmol of each of the oligonucleotides N11 and N12 were mixed in 100  $\mu$ l 12.5 mmol/l Tris-HCl, pH 7.0 and 12.5 mmol/l MgCl<sub>2</sub>, the mixture was heated for 5 min to 95°C and then slowly cooled to room temperature.

N11 : 5' -CATTAAAGAGGAGAAATTAACATGGTATGGTCGACCCCGGGGATATCGC-3'

N12: 5' -TCAGCGATATCCCCGGGGTCTGACCATACCATAGTTAATTTCTCCTCTTTAATGAATT-3'

40

**EcoRI-RBSII-NdeI/Sall/XmaI/EcoRV/CellI linker**

**[0063]**

45

NdeI            SalI    XmaI   EcoRV

5' - CATTAAAGAGGAGAAATTA ACTATGGTATGGTCGACCCGGGGATATCGC - 3'

3'-TTAAGTAATTTCTCCTCTTTAATTGATACCATAACCAGCTGGGGCCCTATAGCGACT-5'

50

EcoRI

Cell III

55

**[0064]** The desired plasmid p16074a was identified by restriction mapping and the expression cassette was checked by DNA sequencing.

**Example 3****Construction of the expression plasmids OripBR-TRP1-EK, OripBR-URA3-EK and OripBR-TET-EK**

5 [0065] The expression cassette (Fig. 3) of the plasmid p16074a composed of the regulated T5-P<sub>N25/03/04</sub> hybrid promoter, the synthetic ribosomal binding site RBSII, a multiple cloning cleavage site with the singular cleavage sites NdeI, Sall, XmaI, EcoRV and CeuI and the  $\lambda$ -T0 terminator was amplified by PCR and inserted into the plasmids OripBR-TRP1, OripBR-URA3 and OripBR-TET.

**3.1 Construction of the expression plasmid OripBR-TRP1-EK**

[0066] The expression cassette (Fig. 3) was amplified by means of PCR using the primers N13 (SEQ ID NO:13) and N14 (SEQ ID NO:14)

15 N13 : 5' -AAAAAAGTAAAGGGGCTTTTCACGG-3'

N14 : 5' -AAAAAAGGATCCGATTCTCACCAATAAAAAACGCCCG-3'

BamHI

20

and the plasmid p16074a as template DNA. A singular BamHI cleavage site was introduced at the 3' end of the expression cassette ( $\lambda$ -T0 terminator) by means of the PCR primer N14.

25 [0067] The PCR product was digested with the restriction endonucleases XhoI and BamHI and the ca. 255 bp long XhoI/BamHI fragment was ligated after purification by agarose gel electrophoresis into the ca. 1490 bp long XhoI/BamHI-OripBR-TRP1 vector fragment (example 2). The desired plasmid OripBR-TRP1-EK (1728 bp) was identified by restriction mapping and the sequence of the expression cassette amplified by PCR was verified by DNA sequencing.

30

**3.2 Construction of the expression plasmid OripBR-URA3-EK**

[0068] The expression plasmid OripBR-URA3-EK (1928 bp) was prepared analogously to the plasmid OripBR-TRP1-EK. However, the base vector OripBR-URA3 was used instead of the base vector OripBR-TRP1.

35

**3.3 Construction of the reference plasmid OripBR-TET-EK**

[0069] The expression cassette (Fig. 3) was amplified by means of PCR using the primers N13 (SEQ ID NO:13) and N14 (SEQ ID NO:15)

40

N15 : 5' -AAAAAATCTAGAGATTCTCACCAATAAAAAACGCCCG-3'

XbaI

45

and the plasmid p16074a as template DNA. A singular XbaI cleavage site was introduced at the 3' end of the expression cassette ( $\lambda$ -T0 terminator) by means of the PCR primer N15.

50 [0070] The PCR product was digested with the restriction endonucleases XhoI and XbaI and the ca. 255 bp long XhoI/XbaI fragment was ligated after purification by agarose gel electrophoresis into the ca. 2000 bp long XhoI/XbaI-OripBR-TET vector fragment (example 2). The desired plasmid OripBR-TET-EK (2243 bp) was identified by restriction mapping and the sequence of the expression cassette amplified by PCR was verified by DNA sequencing.

55

**Example 4****Construction of the expression plasmids OripUC-TRP1-EK, OripUC-URA3-EK and OripUC-TET-EK**

[0071] The medium copy number origin of replication of the plasmid pBR322 was substituted by the high copy number origin of replication of a pUC plasmid in the plasmids OripUC-TRP1-BK, OripUC-URA3-EK and OripUC-TET-EK.

[0072] For this the pUC origin of replication was amplified by means of a PCR using the primers N1 (SEQ ID NO:1) and N2 (SEQ ID NO:2) (example 1) and the plasmid pUC20 as template DNA. The ca. 650 bp long SfiI/NotI-pUC origin of replication fragment digested with SfiI and NotI was ligated after purification by agarose gel electrophoresis into the corresponding vector fragments SfiI/NotI-OripBR-TRP1-BK, SfiI/NotI-OripBR-URA3-BK and SfiI/NotI-OripBR-TBT-BK. The desired plasmids OripUC-TRP1-BK, OripUC-URA3-EK and OripUC-TBT-EK were identified on the basis of their increased plasmid copy number (3-5-fold increase in plasmid yield after standard preparation).

**Example 5****Construction of IFN- $\alpha$ 2b expression plasmids****5.1 Synthesis of the interferon- $\alpha$ 2b model gene**

[0073] The expression vectors were tested with regard to plasmid stability and IFN- $\alpha$ 2b synthesis using an interferon- $\alpha$ 2b gene (Met-IFN- $\alpha$ 2b structural gene without a signal sequence) as a model gene that has to be expressed heterologously.

[0074] The chemically prepared Met-IFN- $\alpha$ 2b structural gene is based on the amino acid sequence stated in the Index Nominum (International Drug Directory 1992/1993) for mature IFN- $\alpha$ 2b. The Met-IFN- $\alpha$ 2b structural gene additionally contains an ATG start codon. The synthetic ribosomal RBSII binding site and a singular EcoRI cleavage site are located upstream at the 5' end of the Met-IFN- $\alpha$ 2b structural gene (Stüber D. et al., 1990). The codons that are preferably used in E. coli (E. coli codon usage) were taken into consideration in the gene design of the Met-IFN- $\alpha$ 2b gene. In addition suitable singular restriction endonuclease cleavage sites were introduced within the coding region and at the ends of the Met-IFN- $\alpha$ 2b structural gene (EcoRI and HindIII) with regard to the construction of variant genes and recloning the Met-IFN- $\alpha$ 2b DNA segment.

[0075] The IFN- $\alpha$ 2b gene (RBSII-Met-IFN- $\alpha$ 2b gene) was prepared by the Genosys Company (Genosys Biotechnologies Inc. Cambridge, England) from oligonucleotides made by chemical synthesis. The double-stranded RBSII-Met-IFN- $\alpha$ 2b gene was assembled by annealing and ligation of the oligonucleotides and subsequently cloned into the EcoRI and HindIII cleavage site of the E. coli standard vector pBluescriptSK(+) from the Stratagene Company (Stratagene GmbH, Heidelberg, Germany). The predetermined DNA sequence of the cloned RBSII-Met-IFN- $\alpha$ 2b structural gene was confirmed by DNA sequencing. Subsequently the ca. 535 bp long EcoRI/HindIII-RBSII-Met-IFN- $\alpha$ 2b segment was recloned into a pDS56/RBSII plasmid digested with EcoRI and HindIII (pDS-IFN). This enables the transfer of the RBSII-Met-IFN- $\alpha$ 2b gene in the form of a ca. 540 bp long EcoRI/CelII-RBSII-Met-IFN- $\alpha$ 2b fragment.

**5.2 Construction of the IFN- $\alpha$ 2b expression plasmids**

[0076] The RBSII-Met-IFN- $\alpha$ 2b gene was isolated from the plasmid pDS-IFN in the form of a ca. 540 bp long EcoRI/CelII-RBSII-Met-IFN- $\alpha$ 2b fragment and subsequently ligated into each of the plasmids OripBR-TRP1-EK, OripBR-URA3-EK, OripBR-TET-EK, OripUC-TRP1-EK, OripUC-URA3-BK and OripUC-TET-EK that were digested with EcoRI and CelII. The desired plasmids OripBR-TRP1-EK-IFN, OripBR-URA3-EK-IFN, OripBR-TET-EK-IFN, OripUC-TRP1-EK-IFN, OripUC-URA3-EK-IFN and OripUC-TET-EK-IFN were identified by restriction mapping.

**Example 6****Construction of non-revertible trpC and pyrF E. coli host strains**

[0077] The desired auxotrophy markers trpC and pyrF were introduced into already available laboratory strains by directed deletion of the complete gene or of a larger gene fragment by means of homologous recombination according to the method of Hamilton C.M. et al. 1989.

## 6.1 Construction of the deletion plasmids

### 6.1.1. Construction of the deletion plasmid pMAK705-trpD/B

[0078] In order to delete the complete chromosomal trpC gene (1355 bp), 700 - 800 bp of the 5' flanking region (trpD) and of the 3' flanking region (trpB) were each cloned by means of PCR and these were ligated into the vector pMAK705 (Hamilton C.M. et al., 1989) in direct fusion (trpD/B) by means of a common cleavage site (NotI) introduced by the PCR primers. The DNA sequence of the tryptophan operon of *E. coli* is given by Yanofsky, C. et al., 1981.

[0079] The 5' flanking region of the trpC gene (trpD) was amplified using the primers N16 (SEQ ID NO:16) and N17 (SEQ ID NO:17)

N16: 5' -AAAAAAGGATCCGGGCTGAAAGTGGCGAAACACGGC-3'

BamHI

N17: 5' -AAAAAAGCGGCCGCTTACCCTCGTGCCGCCAGTGCG-3'

NotI

and chromosomal *E. coli* DNA as a template. A singular BamHI cleavage site was introduced at the 5' end by means of the primer N16 and a singular NotI cleavage site was introduced at the 3' end of the amplified trpD fragment by means of the primer N17.

[0080] The 3' flanking region of the trpC gene (trpB) was amplified using the primers N18 (SEQ ID NO:18) and N19 (SEQ ID NO:19)

N18: 5' -AAAAAAGCGGCCGCGGAAAGGAACAATGACAACATTACTTAAC-3'

NotI

N19: 5' -AAAAAAGCATGCCGGTGCCCGTGCTCGCCAGTTTCG-3'

SpHI

and chromosomal *E. coli* DNA as a template. A singular NotI cleavage site was introduced at the 5' end by means of the primer N18 and a singular SpHI cleavage site was introduced at the 3' end of the amplified trpB fragment by means of the primer N19.

[0081] The trpD PCR fragment was digested with BamHI and NotI (ca. 710 bp) and the trpB PCR fragment was digested with NotI and SpHI (ca. 820 bp). After purification by agarose gel electrophoresis, the DNA fragment was ligated into the ca. 5650 bp long BamHI/SpHI-pMAK705 vector fragment by means of a three fragment ligation in a direct fusion via the common NotI cleavage site. The desired plasmid pMAK705-trpD/B (7104 bp) was identified by restriction mapping and the trpD/B sequence amplified by PCR was verified by DNA sequencing.

### 6.1.2 Construction of the deletion plasmid pMAK705-ΔpyrF

[0082] In order to delete ca. 260 bp from the coding region of the chromosomal pyrF gene, 700 - 800 bp of the flanking pyrF region was cloned in each case by means of PCR and ligated in a direct fusion (ΔpyrF) into the vector pMAK705 by means of a common cleavage site (XbaI) introduced by the PCR primers. The DNA sequence of the pyrF operon of *E. coli* was published by Turnbough, C.L.Jr. et al., 1987. The 5' flanking region of the ΔpyrF gene was amplified using the primers N20 (SEQ ID NO:20) and N21 (SEQ ID NO:21)

N20: 5' AAAAAAGGATCCGTACGTAGTAAGCCTCGTTATCGTTG-3'

BamHI

N21: 5'-AAAAAATCTAGACACACGCCTAAGTCAGCTGCAGC-3'

XbaI

and chromosomal E. coli DNA as a template. A singular BamHI cleavage site was introduced at the 5' end by means of the primer N20 and a singular XbaI cleavage site was introduced at the 3' end of the amplified DNA fragment by means of the primer N21.

[0083] The 3' flanking region of the  $\Delta$ pyrF gene was amplified using the primers N22 (SEQ ID NO:22) and N23 (SEQ ID NO:23)

N22: 5' -AAAAAATCTAGAGGTCAGGAGTTCAAACTGGTTACG-3'

XbaI

N23: 5' -AAAAAAGCATGCCAGCTGTCGGAGGTGTTATTGGTC-3'

SpHI

and chromosomal E. coli DNA as a template. A singular XbaI cleavage site was introduced at the 5' end by means of the primer N22 and a singular SpHI cleavage site was introduced at the 3' end of the amplified DNA fragment by means of the primer N23.

[0084] The 5' flanking  $\Delta$ pyrF fragment was digested with BamHI and XbaI (ca. 600 bp) and the 3' flanking  $\Delta$ pyrF fragment was digested with XbaI and SpHI (ca. 690 bp). After purification by agarose gel electrophoresis, the DNA fragments were ligated in a direct fusion into the ca. 5650 bp long BamHI/SpHI-pMAK705 vector fragment by means of a three fragment ligation via the common XbaI cleavage site. The desired plasmid pMAK705- $\Delta$ pyrF (6788 bp) was identified by restriction mapping and the pyrF sequence amplified by PCR was verified by DNA sequencing.

## 6.2 Isolation of E. coli deletion mutants

[0085] The vector pMAK705 is characterized by a temperature-sensitive origin of replication (origin of replication from pSC101) which is inactive at 44°C and by a chloramphenicol resistance gene. In order to delete a chromosomal gene the appropriate pMAK705 plasmid derivative is transformed in the desired E. coli strain. A prerequisite for a homologous recombination between the plasmid and E. coli chromosome is a recombination-competent E. coli strain (recA<sup>+</sup> genotype). Strains whose genotype is recA<sup>-</sup> cannot carry out a homologous recombination and are therefore unsuitable.

[0086] The E. coli K12 strain UT5600 (ara, proC-14, leuB6, azi-6, lacY1, tsx-67, entA403, trpE38<sup>+</sup>, rpsL109, xyl-5, mtl-1, thi-1,  $\Delta$ ompT) (Grodberg, J. and Dunn, J.J., 1988) was transformed with the plasmid pMAK705-trpD/B and with the plasmid pMAK705- $\Delta$ pyrF. After addition of the plasmid DNA, the transformation preparation was incubated for 30 minutes on ice, for 5 minutes at 37°C (heat shock) and for 30 minutes at 30°C. The transformation preparation was transferred to 3 ml LB medium containing 20  $\mu$ g/ml chloramphenicol and the cells were cultured overnight.

[0087] In order to select for plasmid cointegrates, i.e. for cells which have integrated the plasmid into the chromosome by means of homologous recombination, 300  $\mu$ l of a 10<sup>-5</sup> dilution of the overnight culture was plated out on LB agar plates containing 20  $\mu$ g/ml chloramphenicol. The agar plates were preheated to 50 - 60°C in an incubator and, after plating out the cells, were immediately incubated at exactly 44°C. Since the pMAK705 plasmid does not replicate at this temperature, only those cells grow into colonies which have chromosomally integrated the plasmid. The cointegrate candidates were subsequently passaged 2 to 3 times on LB agar plates containing 20  $\mu$ g/ml chloramphenicol at 44°C for purification (dilution smear).

[0088] In order to break up the cointegrates and to cure the plasmid that disintegrated from the chromosome, a purified cointegrate clone was cultured for several passages (2 to 6) in LB medium at 30°C in the absence of chloramphenicol. Afterwards the cells were separated by dilution on LB medium and tested for the desired genotype.

### 6.2.1. Characterization of the deletion mutant UT5600 ( $\Delta$ trpC)

[0089] The cells/clones separated by dilution on LB medium were analysed with regard to the desired trp auxotrophy by plating out on medium (M9 minimal medium containing 0.5 % casamino acids) with tryptophan (50 mg/ml) and without tryptophan (replica plating). In addition the chloramphenicol sensitivity of the clones was checked by plating out on LB medium with chloramphenicol (20  $\mu$ g/ml) and without chloramphenicol. In addition the chromosomal trpC deletion was confirmed by comparative PCR analysis between the initial strain and deletion mutant using the PCR primers N1 and N4 and the corresponding chromosomal DNA as template DNA. The PCR product of the initial strain was ca. 2900 bp long as calculated theoretically and that of the trpC deletion mutant was ca. 1430 bp long.

### 6.2.2. Characterization of the deletion mutant UT5600 ( $\Delta$ pyrF)

[0090] The cells/clones separated by dilution on LB medium were analysed with regard to the desired pyrF auxotrophy by plating out on medium (M9 minimal medium containing 0.5 % casamino acids) with uracil (20 mg/ml) and without uracil. In addition the chloramphenicol sensitivity of the clones was checked by plating out on LB medium with chloramphenicol (20  $\mu$ g/ml) and without chloramphenicol. In addition the chromosomal pyrF deletion was confirmed by comparative PCR analysis between the initial strain and deletion mutant using the PCR primers N5 and N8 and the corresponding chromosomal DNA as template DNA. The PCR product of the initial strain was ca. 1550 bp long as calculated theoretically and that of the pyrF deletion mutant was ca. 1290 bp long.

## Example 7

### Expression of a heterologous model gene (IFN- $\alpha$ 2b)

[0091] The new host/vector system based on antibiotic-free selection by complementation of an auxotrophy was evaluated with regard to its function and performance using interferon- $\alpha$  2b as a model gene that has to be expressed heterologously.

#### 7.1 Selection by complementation of the trpC auxotrophy

[0092] In order to express the IFN- $\alpha$ 2b gene, the E. coli K12 strain UT5600 ( $\Delta$ trpC) (example 6) was transformed in each case with one of the expression plasmids OripBR-TRP1-EK-IFN, OripUC-TRP1-EK-IFN, OripBR-TET-EK-IFN and OripUC-TET-EK-IFN described in example 5. The transformed UT5600( $\Delta$ trpC)/OripBR-TRP1-EK-IFN and UT5600 ( $\Delta$  trpC)/OripUC-TRP1-EK-IFN cells were grown at 37°C in a shaking culture in M9 minimal medium containing 0.5 % casamino acids (Difco) and the reference cells UT5600 ( $\Delta$ trpC)/OripBR-TET-EK-IFN and OripUC-TET-EK-IFN were grown in M9 minimal medium containing 0.5 % casamino acids (Difco) and 50 mg/ml tryptophan and 12.5  $\mu$ g/ml tetracycline at 37°C up to an optical density at 550 nm ( $OD_{550}$ ) of 0.6 - 0.9 and subsequently induced with IPTG (1 - 5 mmol/l final concentration). After an induction phase of 4 - 8 hours (h) at 37°C, the cells were harvested by centrifugation (Sorvall RC-5B centrifuge, GS3 rotor, 6000 rpm, 15 min), washed with 50 mmol/l Tris-HCl buffer, pH 7.2 and stored at -20°C until further processing.

#### 7.2 Selection by complementation of the pyrF auxotrophy

[0093] In order to express the IFN- $\alpha$ 2b gene, the E. coli K12 strain UT5600 ( $\Delta$ pyrF) (example 6) was transformed in each case with one of the expression plasmids OripBR-URA3-EK-IFN, OripUC-URA3-EK-IFN, OripBR-TET-EK-IFN and OripUC-TET-EK-IFN described in example 5. The transformed UT5600( $\Delta$ trpC)/OripBR-URA3-EK-IFN and UT5600 ( $\Delta$  trpC)/OripUC-URA3-EK-IFN cells were grown at 37°C in a shaking culture in M9 minimal medium containing 0.5 % casamino acids (Difco) and the reference cells UT5600 ( $\Delta$ trpC)/OripBR-TET-EK-IFN and OripUC-TET-EK-IFN were grown in M9 minimal medium containing 0.5 % casamino acids (Difco) and 20 mg/ml uracil and 12.5  $\mu$ g/ml tetracycline at 37°C up to an optical density at 550 nm ( $OD_{550}$ ) of 0.6 - 0.9 and subsequently induced with IPTG (1 - 5 mmol/l final concentration). After an induction phase of 4 - 8 hours (h) at 37°C, the cells were harvested by centrifugation (Sorvall RC-5B centrifuge, GS3 rotor, 6000 rpm, 15 min), washed with 50 mmol/l Tris-HCl buffer, pH 7.2 and stored at -20°C until further processing.

#### 7.3 Expression analysis in the standard system

[0094] The synthesis of IFN- $\alpha$ 2b was analysed in the UT5600 ( $\Delta$ trpC) or UT5600 ( $\Delta$ pyrF) cells transformed with the expression plasmids OripBR-TRP1-EK-IFN, OripUC-TRP1-EK-IFN, OripBR-URA3-EK-IFN, OripUC-URA3-EK-IFN,

OripBR-TET-EK-IFN and OripUC-TET-EK-IFN. For this cell pellets from 3 OD<sub>550</sub> units (1 OD<sub>550</sub> = 1 ml cell suspension with an OD<sub>550</sub> of 1) of centrifuged culture medium was resuspended in 0.25 ml 10 mmol/l Tris-HCl, pH 7.2 and the cells were lysed by ultrasonic treatment (2 pulses of 30 s at 50 % intensity) with a Sonifier® Cell Disruptor B15 from the Branson Company (Heusenstamm, Germany). The insoluble cell components were sedimented (Eppendorf 5415 centrifuge, 14000 rpm, 5 min) and the supernatant was admixed with 1/5 volumes (vol) 5xSDS sample buffer (1xSDS sample buffer: 50 mmol/l Tris-HCl, pH 6.8, 1 % SDS, 1 % mercaptoethanol, 10 % glycerol, 0.001 % bromophenol blue). The insoluble cell debris fraction (pellet) was resuspended in 0.3 ml 1xSDS sample buffer containing 6-8 M urea, the samples were incubated for 5 min at 95°C and again centrifuged. Subsequently the proteins were separated by SDS polyacrylamide gel electrophoresis (PAGE) (Laemmli, U.K., 1970) and stained with Coomassie Brilliant Blue R dye.

[0095] The synthesized IFN- $\alpha$ 2b protein was homogeneous and was found exclusively in the insoluble cell debris fraction in the form of insoluble protein aggregates, the so-called inclusion bodies (IBs). The expression yield was comparable within the scope of the measurement accuracy in all clones/cells and was between 30 - 50 % relative to the total E. coli protein.

### Example 8

#### Examination of the plasmid stability

[0096] A sample was taken before induction of the cells (example 7) and a plasmid stability test was carried out. For this the samples were diluted directly with M9 minimal medium containing 0.5 % casamino acids and 300  $\mu$ l of each dilution step was plated out on a non-selective agar plate (M9 minimal medium supplemented with 0.5 % casamino acids, 50 mg/ml tryptophan and 20 mg/l uracil). Subsequently 400 individual colonies from each clone were firstly smeared with a toothpick on an appropriate selective agar plate (M9 minimal medium containing 0.5 % casamino acids or M9 minimal medium supplemented with 0.5 % casamino acids, 50 mg/l tryptophan, 20 mg/ml uracil and 12.5  $\mu$ g/ml tetracycline) and then on a non-selective agar plate (M9 minimal medium supplemented with 0.5 % casamino acids, 50 mg/l tryptophan and 20 mg/l uracil).

[0097] The number of plasmid-containing cells was determined from the number of cells which formed a colony on both agar plates. Cells which do not grow on the selective agar plate have lost the plasmid for complementing the auxotrophy or forming the antibiotic-resistance (TET reference plasmids). Before induction of the cells the plasmid stability was 100 % for all cultures independent of the selection method and independent of the plasmid type.

#### List of references

##### [0098]

- Amann, E., et al., Gene 25 (1983) 167-178
- Bachman, B.J., Microbiol. Rev. 47 (~983) 180-230
- Backman, K. et al. Cold Spring Harbor Symposia Quant. Biol. 43 (1978) 69-76
- Beck, F. et al., Gene 16 (1981) 35-58
- Boliver, F. et al., Gene 2 (1979) 95-113
- Botstein, D., et al., Gene 8 (1979) 17-24
- Brosius, J., et al., J. Mol. Biol. 148 (1981) 107-127
- Bujard, H., et al., Methods in Enzymology 155 (1987) 416-433
- Carter, P. et al., Biotechnology 10 (1992) 163-167
- Chambers, S.P., et al., Appl. Microbiol. Biotechnol. 29 (1988) 572-578
- DeBoer, H.A., et al., Proc. Natl. Acad. Sci. USA 80 (1983) 21-25.
- EP-A 0651803
- Fiedler, S., et al., Anal. Biochem. 170 (1988) 38-44
- Grodberg, J., et al., J. Bacteriol. 170 (1988) 1245-1253
- Hamilton, C.M., et al., J. Bacteriol. 171 (1989) 4617-4622
- Hanahan, D., J. Mol. Biol. 166 (1983) 557-580
- Harley, C.B., Nucl. Acids Res. 15 (1987) 2343-2361
- Higgins, C.F.; Causton H.C.; Dance, G.S.C.; Mudd, E.A.: The role of 3' end in mRNA stability and decay. In: Belasco, J.G. Brawerman (eds.). Control of Messenger RNA Stability. Academic Press, Inc., San Diego, CA, p.13-30 (1993)
- Index Nominum. International Drug Directory 1992/1993, Medpharm Scientific Publishers & Wissenschaftliche Verlagsgemeinschaft Stuttgart, Govi-Verlag Frankfurt, 615-616 (1992)
- Kopetzki, E., et al., Mol. Gen. Genet. 216 (1989) 149-155



Kopetzki, E., et al., Yeast 5 (1989) 11-24

Laemmli, U.K., et al., Nature 227 (1970) 680-685

Miller, J.H.: Experiments in Molecular Genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1972)

Mullis, K.B. et al., Methods Enzymol. 155 (1987) 350-355

Neidhardt, F.C.; Ingraham, J.L.; Low, K.B.; Magasanik, B.; Schaechter, M.; Umberger, H.E.: Escherichia coli and Salmonella typhimurium. Cellular and Molecular Biology. American Society for Microbiology, Washington D.C. (1987)

Rose, M., et al., Gene 29 (1984) 113-124

Rosenberg, M., et al., Ann. Rev. Genet. 13 (1979) 319-353

Sambrook, J.; Fritsch, E.F.; Maniatis, T.: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, Cold Spring Harbor, New York (1989)

Schlegel, H.G.: Allgemeine Mikrobiologie. Georg Thieme Verlag Stuttgart, New York (1981)

Schwarz, E., et al., Nature 272 (1978) 410-414

Struhl, K., et al., Proc.Natl.Acad.Sci. USA 73 (1976) 1471-1475

Stüber, D., et al., Immunological Methods IV (1990) 121-152

Studier, F., et al., Methods Enzymol. 185 (1990) 60-89

Sutcliffe, G., et al., Quant. Biol. 43 (1979) 77-90

Tschumper, G., Gene 10 (1980) 157-166

Turnbough, C.L.Jr., et al., J.Biol.Chem. 262 (1987) 10239-10245

Weir, A.N.C. and Mountain, A. EP 0 651 1803

Winnacker, E.L., Gene und Klone, VCH Weinheim, DE, 1985

Yanofsky, C., et al., Nucl.Acids Res. (1981) 6647-6668

# EP 0 972 838 A1

## SEQUENCE LISTING

5 <110> Roche Diagnostics GmbH  
 <120> New Escherichia coli host/vector system based on  
 antibiotic-free selection by complementation of an  
 auxotrophy  
 <130> Case 20315  
 10 <140>  
 <141>  
 <150> EP98113156.8  
 <151> 1998-07-15  
 15 <150> EP98119078.8  
 <151> 1998-10-09  
 <160> 23  
 <170> PatentIn Ver. 2.0  
 20 <210> 1  
 <211> 41  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 25 <223> Description of Artificial Sequence:primer  
 <400> 1  
 aaaaaaggcc atataggccg gccgcgttgc tggcgttttt c 41  
 <210> 2  
 <211> 35  
 <212> DNA  
 30 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence:primer  
 <400> 2  
 35 aaaaaactcg aggagcgtca gaccccgtag aaaag 35  
 <210> 3  
 <211> 53  
 <212> DNA  
 40 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence:primer  
 <400> 3  
 45 aaaaaaggcc atataggcca tatgcggccg cggagagggc caagagggag ggc 53  
 <210> 4  
 <211> 104  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 50 <223> Description of Artificial Sequence:primer  
 <400> 4  
 55

# EP 0 972 838 A1

aaaaaactcg agatatggat cctctccaaa aaaaaaggct ccaaaggag cctttaattg 60  
 tatcggttta ttactatatt cttagcattt ttgacgaaat ttgc 104

5 <210> 5  
 <211> 58  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 10 <223> Description of Artificial Sequence:primer

<400> 5  
 aaaaaaggcc atataggcca tatgcgcccg ccggtaatct ccgaacagaa ggaagaac 58

<210> 6  
 <211> 102  
 15 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer

<400> 6  
 20 aaaaaactcg agacgaggat ccataaacg atacaattaa aggctccttt tggagccttt 60  
 ttttttgag aactttagtt ttgctggccg catcttctca aa 102

<210> 7  
 <211> 60  
 25 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer

<400> 7  
 30 aaaaaaggcc atataggcca tatgcgcccg cctcatgttt gacagcttat catcgataag 60

<210> 8  
 <211> 94  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> Description of Artificial Sequence:primer

<400> 8  
 40 aaaaaactcg agatattcta gaataaacg atacaattaa aggctccttt tggagccttt 60  
 ttttttgag atcaggtcga ggtggcccg ctcc 94

<210> 9  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <223> Description of Artificial Sequence:primer

<400> 9  
 aggccggggg ggggc 15

50 <210> 10  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

55

**EP 0 972 838 A1**

	<220>	
	<223> Description of Artificial Sequence:primer	
5	<400> 10	
	ggccgcccc ccccgcccta ta	22
	<210> 11	
	<211> 50	
	<212> DNA	
10	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence:primer	
	<400> 11	
15	cattaaagag gagaaattaa ctatggtatg gtcgaccccg gggatatcgc	50
	<210> 12	
	<211> 57	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Description of Artificial Sequence:primer	
	<400> 12	
	tcagcgatat ccccggggtc gaccatacca tagttaattt ctctcttcta atgaatt	57
25	<210> 13	
	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence:primer	
30	<400> 13	
	aaaaaaagtaa aggggctttt cacgg	25
	<210> 14	
	<211> 37	
35	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence:primer	
	<400> 14	
40	aaaaaaggat ccgattctca ccaataaaaa acgccccg	37
	<210> 15	
	<211> 37	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Description of Artificial Sequence:primer	
	<400> 15	
	aaaaaatcta gagattctca ccaataaaaa acgccccg	37
50	<210> 16	
	<211> 36	
	<212> DNA	
	<213> Artificial Sequence	
55		

<220>  
 <223> Description of Artificial Sequence:primer  
 5 <400> 16  
 aaaaaaggat ccgggctgaa agtggcgaaa cacggc 36  
 <210> 17  
 <211> 36  
 <212> DNA  
 10 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence:primer  
 15 <400> 17  
 aaaaaagcgg ccgcttacct tcgtgccgcc agtgcg 36  
 <210> 18  
 <211> 43  
 <212> DNA  
 20 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence:primer  
 <400> 18  
 25 aaaaaagcgg ccgcggaaaag gaacaatgac aacattactt aac 43  
 <210> 19  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Description of Artificial Sequence:primer  
 <400> 19  
 aaaaaagcat gccggtgcgc cgtgctcgcc agtttcg 37  
 35 <210> 20  
 <211> 38  
 <212> DNA  
 <213> Artificial Sequence  
 40 <220>  
 <223> Description of Artificial Sequence:primer  
 <400> 20  
 aaaaaaggat ccgtacgtag taagcctcgt ttcggtg 38  
 45 <210> 21  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence:primer  
 50 <400> 21  
 aaaaaatcta gacacagcc taagtcagct gcagc 35  
 55

<210> 22  
 <211> 36  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer

<400> 22  
 aaaaaatcta gaggtcagga gttcaaactg gttacg 36

<210> 23  
 <211> 36  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer

<400> 23  
 aaaaaagcat gccagctgtc ggaggtgtta ttggtc 36

## Claims

1. Minimal prokaryotic expression vector which cannot be homologously recombined with the genome of prokaryotic organisms containing:
  - a) an origin of replication,
  - b) an auxotrophy marker gene,
  - c) a promoter which is functionally active in prokaryotes and
  - d) a foreign gene to be expressed which is under the control of the said promoter.
2. Expression vector as claimed in claim 1, wherein the auxotrophy marker gene and the gene to be expressed are oriented in opposite directions and are terminated by the same terminator.
3. Expression vector as claimed in claims 1 to 2, wherein the auxotrophy marker is derived from eukaryotes.
4. Expression vector as claimed in claims 1 to 3, wherein the auxotrophy marker compensates a tryptophan, leucine or uracil deficiency.
5. Expression vector as claimed in claims 1 to 4, wherein the expression yield of the auxotrophy marker gene is 1 % or less of the total protein in the host cell when the auxotrophy is complemented in the said host cell.
6. Prokaryotic host cell containing a vector as claimed in claims 1 to 5, wherein the host cell contains a mutation in a chromosomal gene which corresponds to the auxotrophy marker gene of the expression vector and the result of the mutation is that no functional product of the said chromosomal host cell gene can be expressed.
7. Process for the construction of a prokaryotic expression system, wherein an expression vector as claimed in claims 1 to 5 is introduced into a prokaryotic host cell with a mutation in a chromosomal gene which is complemented by the auxotrophy marker gene of the expression vector and the result of the chromosomal mutation is that no function product of the said chromosomal host cell gene can be expressed.
8. Process for the production of a heterologous protein by heterologous expression of a foreign gene in an expression

vector as claimed in claims 1 to 5 which is contained in a host cell as claimed in claim 6 and isolation of the expression product from the host cell or the supernatant, wherein the fermentation of the host cell is carried out in a minimal medium or a synthetic complete medium.

5

10

15

20

25

30

35


40

45

50

55

Fig. 1








Fig. 3

```

XhoI
ctcgaagAAATCATAAAAAATTTATTTGCTTTGTGAGCGGATAACAATTATAATAGATTCA 60
|--- T5-Pu25/03/04 --->

EcoRI NdeI SalI
ATTGTGAGCGGATAACAATTTACACAGAATTCATTAAAGAGGAGAAATTACAATATGgtc 120
RBSII

XmaI EcoRV CclII
gacccccggggaatcGCTGAGCTTGGACTCCTGTTGATAGATCCAGTAATGACCTCAGAA 180
|--- λ -T0 Terminator --->

CTCCATCTGGATTTGTTTCAGAACGCTCGGTTGCCGCCGGGCGTTTTTTTATTGGTGAGAAT 240

BamHI
Cggatcc 247
|

```



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 99 11 3182

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	EP 0 178 764 A (GENEX CORP) 23 April 1986 (1986-04-23) * the whole document *	1,4,6-8	C12N15/70 C12N15/64 C12N15/65 C07K14/56 //C12N15/21
X	EP 0 185 512 A (NOVO INDUSTRI AS) 25 June 1986 (1986-06-25) * the whole document *	1,6-8	
Y	---	2-5	
X	FR 2 511 032 A (CENTRE NAT RECH SCIENT) 11 February 1983 (1983-02-11) * the whole document *	1,6-8	
Y	---	2-5	
X	EP 0 258 118 A (TRANSGENE SA) 2 March 1988 (1988-03-02) * the whole document *	1,6-8	
Y	---	2-5	
X	WO 89 03427 A (UNIV WASHINGTON) 20 April 1989 (1989-04-20) * the whole document *	1,6-8	
Y	---	2-5	
X	E. DEGRYSE: "Stability of a host-vector system based on complementation of an essential gene in Escherichia coli" J. BIOTECHNOLOGY, vol. 18, no. 1-2, April 1991 (1991-04), pages 29-40, XP002092102 ELSEVIER, AMSTERDAM,NL * the whole document *	1,6-8	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N C07K
Y	---	2-5	
X	EP 0 157 441 A (UNILEVER NV) 9 October 1985 (1985-10-09) * page 2, line 14 - line 35 * * page 9, line 11 - line 20; claims 1,4; figures 1,8 *	1,6-8	
Y	---	2-5	
	--- -/--		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 6 October 1999	Examiner Hornig, H
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (F04C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 99 11 3182

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	J.E. HILL ET AL.: "Yeast/E.coli shuttle vectors with multiple unique restriction sites" YEAST, vol. 2, no. 3, September 1986 (1986-09), pages 163-167, XP002092103 John Wiley & Sons Ltd. , Chichester, UK * the whole document *	1-8	
D,Y	M. ROSE ET AL.: "Structure and function of the yeast URA3 gene: expression in Escherichia coli" GENE, vol. 29, 1984, pages 113-124, XP002092104 ELSEVIER SCIENCE PUBLISHERS,B.V.,AMSTERDAM,NL; * the whole document *	1-8	
D,Y	D. BOTSTEIN ET AL.: "Sterile host yeast (SHY): A eucaryotic system of biological containment for recombinant DNA experiments" GENE, vol. 8, no. 1, December 1979 (1979-12), pages 17-24, XP002092105 ELSEVIER SCIENCE PUBLISHERS,B.V.,AMSTERDAM,NL; * page 20, line 15 - line 22 *	1-8	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 6 October 1999	Examiner Hornig, H
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P4/C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 99 11 3182

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	R.D. GIETZ AND A. SUGINO: "New yeast-Escherichia coli shuttle vector constructed with in vitro mutagenized yeast genes lacking sic-base pair restriction site" GENE, vol. 74, 1988, pages 527-534, XP002092106 ELSEVIER SCIENCE PUBLISHERS, B.V., AMSTERDAM, NL; * page 531, left-hand column, line 11 - line 20 *	1-8	
Y	WO 91 13971 A (HAWAII BIOTECH GROUP) 19 September 1991 (1991-09-19) * page 5, line 15 - line 22 *	1-8	
D, A	WO 94 02607 A (CELLTECH LTD ; WEIR ANDREW NEIL CHARLES (GB); MOUNTAIN ANDREW (GB)) 3 February 1994 (1994-02-03) * the whole document *	1-8	
A	WO 91 11523 A (UPJOHN CO) 8 August 1991 (1991-08-08) * the whole document *	1-8	
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>6 October 1999</b>	Examiner <b>Hornig, H</b>
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 11 3182

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-10-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0178764 A	23-04-1986	JP 61058595 A	25-03-1986
EP 0185512 A	25-06-1986	DK 594084 A	13-06-1986
		AT 55413 T	15-08-1990
		AU 592542 B	18-01-1990
		AU 5108185 A	19-06-1986
		CA 1292197 A	19-11-1991
		DK 571985 A,B,	13-06-1986
		FI 854893 A,B,	13-06-1986
		JP 2571338 B	16-01-1997
		JP 6086669 A	29-03-1994
		JP 2576970 B	29-01-1997
		JP 61139392 A	26-06-1986
		US 4920048 A	24-04-1990
FR 2511032 A	11-02-1983	NONE	
EP 0258118 A	02-03-1988	FR 2602792 A	19-02-1988
		FR 2618159 A	20-01-1989
		AT 129285 T	15-11-1995
		AU 616637 B	07-11-1991
		AU 7658987 A	11-02-1988
		DE 3751564 D	23-11-1995
		DE 3751564 T	28-03-1996
		DK 408687 A	06-02-1988
		ES 2079348 T	16-01-1996
		GR 3018296 T	31-03-1996
		JP 2905921 B	14-06-1999
		JP 10004981 A	13-01-1998
		JP 2772793 B	09-07-1998
		JP 63233790 A	29-09-1988
		JP 2844191 B	06-01-1999
		JP 10066575 A	10-03-1998
		US 5198343 A	30-03-1993
WO 8903427 A	20-04-1989	AT 121785 T	15-05-1995
		AU 637969 B	17-06-1993
		AU 2611188 A	02-05-1989
		DE 3853683 D	01-06-1995
		DE 3853683 T	31-08-1995
		EP 0381706 A	16-08-1990
		JP 4501351 T	12-03-1992
		US 5387744 A	07-02-1995
		US 5855879 A	05-01-1999
		US 5672345 A	30-09-1997
		US 5840483 A	24-11-1998

EPO FORM P4459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 11 3182

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-10-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 8903427 A		US 5855880 A	05-01-1999
		US 5294441 A	15-03-1994
EP 0157441 A	09-10-1985	NL 8400687 A	01-10-1985
		AT 72832 T	15-03-1992
		DE 3585415 A	02-04-1992
		DK 504585 A	01-11-1985
		WO 8503945 A	12-09-1985
		JP 2711258 B	10-02-1998
		JP 62500280 T	05-02-1987
		US 5683909 A	04-11-1997
WO 9113971 A	19-09-1991	AU 7493491 A	10-10-1991
		US 5695965 A	09-12-1997
WO 9402607 A	03-02-1994	DE 69310373 D	05-06-1997
		DE 69310373 T	25-09-1997
		EP 0651803 A	10-05-1995
		US 5928903 A	27-07-1999
WO 9111523 A	08-08-1991	AT 116687 T	15-01-1995
		AU 4181593 A	30-09-1993
		AU 643752 B	25-11-1993
		AU 7184091 A	21-08-1991
		CA 2070994 A	01-08-1991
		DE 69106498 D	16-02-1995
		DE 69106498 T	24-05-1995
		DK 514405 T	10-07-1995
		EP 0514405 A	25-11-1992
		ES 2069277 T	01-05-1995
		GR 3015629 T	30-06-1995
		JP 5504063 T	01-07-1993

EPO FORM P469

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82